IJISET - International Journal of Innovative Science, Engineering & Technology, Vol. 09 Issue 08, August 2022
ISSN (Online) 2348 – 7968 | Impact Factor – 6.72
www.ijiset.com

On the Hypersurface of a Finsler space with generalized Matsumoto metric

Surekha Desai[1], Narasimhamurthy S K[2] and Raghavendra R S[3]

Department of P.G. Studies and Research in Mathematics, Kuvempu University,
Shankaraghatta-577 451, Shivamogga, Karnataka, INDIA.
e-mail: surekhad90@gmail.com, nmurthysk@hotmail.com and rsr.smg@gmail.com

Abstract

In this paper, we study some geometrical properties of a Finsler space with the generalized Matsumoto metric $L = \frac{\alpha^{m+1}}{(\alpha-\beta)^m}$. Further, we prove the necessary and sufficient condition for Finsler Hypersurface satisfies the condition of hyperplanes of first, second kinds and but not the hyperplane of the third kind with respect to the above metric.

Keywords: Finsler space, Generalized Matsumoto metric, Hyperplanes, Induced Cartan's connection, $(\boldsymbol{\alpha}, \boldsymbol{\beta})$-metric.

1. Introduction

In 1992,[1] the notion of Finsler spaces with an (α, β)-metric was first proposed by M. Matsumoto named as function of $L(\alpha, \beta)$. After the Matsumoto's accomplishment in the development of Finsler geometry there are lot of contributions were given by several authors they have studied a special form of (α, β)-metrics like Rander's metric, Kropina metric, generalized Kropina metric, Shen's square metric etc. The systematic theory of the hypersurface of a Finsler space was built by Matsumoto in 1985, along with this he explained the hyperplane of the first kind, second kind and third kind are the classification of hypersurfaces. Further, many researchers were considered these three kinds of hyperplanes in different types of (α, β)-metrics of Finsler spaces and they came with various conclusions. Recent years, in 2009, H. G. Nagaraja, S. K. Narasimhamurthy, Pradeep Kumar and S. T. Aveesh obtained some results on geometrical properties of Finslerian hypersurfaces with (α, β)-metrics [3, 4]. In 2018, K. Vineet and R. K. Gupta worked on some special (α, β)-metric. In 2020, Brijesh kumar Tripathi introduced same aspect with deformed Berwald-infinite series metric.

In this paper, our aim is to express certain geometrical properties of hypersurface of a Finsler spaces, and we discussed the different kinds of hyperplanes with generalized Matsumoto metric, $L = \frac{\alpha^{m+1}}{(\alpha-\beta)^m}$. We have derived that the necessary and sufficient condition for hypersurface of generalized Matsumoto metric satisfies the conditions of hyperplane of first, second and but not third kind for above metric.

IJISET - International Journal of Innovative Science, Engineering & Technology, Vol. 09 Issue 08, August 2022
ISSN (Online) 2348 – 7968 | Impact Factor – 6.72
www.ijiset.com

2. Preliminaries

We consider an n-dimensional Finsler space with smooth manifold M^n assigned with L i.e., $F_n = \left(M^n,\ L(\alpha,\beta)\right)$, where α-Riemannian metric and β-differential 1-form. Here

$$L = \frac{\alpha^{m+1}}{(\alpha-\beta)^m}. \tag{2.1}$$

Now differentiate the Eq. (2.1) partially with respect to α and β for first and second order, we get,

$$\left.\begin{aligned} L_\alpha = \frac{\alpha^m(\alpha-m\beta-\beta)}{(\alpha-\beta)^{m+1}}, L_\beta = \frac{m\alpha^{m+1}}{(\alpha-\beta)^{m+1}}, \\ L_{\alpha\alpha} = -2m(m+1)\left(\frac{\alpha^m}{(\alpha-\beta)^{m+1}} - \frac{\alpha^{m-1}}{2(\alpha-\beta)^m} - \frac{\alpha^{m+1}}{2(\alpha-\beta)^{m+2}}\right), \\ L_{\beta\beta} = \frac{m(m+1)\alpha^{m+1}}{(\alpha-\beta)^{m+2}},\ L_{\alpha\beta} = m(m+1)\left(\frac{\alpha^m}{(\alpha-\beta)^{m+1}} - \frac{\alpha^{m+1}}{(\alpha-\beta)^{m+2}}\right) \end{aligned}\right\} \tag{2.2}$$

where, $L_\alpha = \frac{\partial L}{\partial \alpha}, L_\beta = \frac{\partial L}{\partial \beta}, L_{\alpha\alpha} = \frac{\partial L_\alpha}{\partial \alpha}, L_{\beta\beta} = \frac{\partial L_\beta}{\partial \beta}, L_{\alpha\beta} = \frac{\partial L_\alpha}{\partial \beta}$.

The normalized element with supporting element $l_i = \dot{\partial}_i L$ and angular metric tensor $h_{ij} = L^{-1}\dot{\partial}_i\dot{\partial}_j L$ is given by,

$$\left.\begin{aligned} l_i = \alpha^{-1} L_\alpha Y_i + L_\beta b_i \\ h_{ij} = P a_{ij} + Q_0\, b_i\, b_j + Q_1\left(b_i\, Y_j + b_j\, Y_i\right) + Q_2\, Y_i\, Y_j \end{aligned}\right\} \tag{2.3}$$

where, $Y_i = a_{ij}y^j, \dot{\partial}_i = \frac{\partial}{\partial y^i}$ and

$$P = LL_\alpha\alpha^{-1}, \qquad Q_0 = LL_{\beta\beta}, \qquad Q_1 = LL_{\alpha\beta}\alpha^{-1}, \qquad Q_2 = L\alpha^{-2}(L_{\alpha\alpha} - L_\alpha\alpha^{-1}) \tag{2.4}$$

The constant quantities of fundamental function $L(\alpha,\beta)$ of Eq. (2.1) are given below by using Eq. (2.4), we get

$$\left.\begin{aligned} P = \frac{\alpha^{2m}(m+1)}{(\alpha-\beta)^{2m}} - \frac{m\alpha^{2m+1}}{(\alpha-\beta)^{2m+1}}, \\ Q_0 = \frac{m(m+1)\alpha^{2m+2}}{(\alpha-\beta)^{2m+2}}, \\ Q_1 = m(m+1)\left(\frac{\alpha^{2m}}{(\alpha-\beta)^{2m+1}} - \frac{\alpha^{2m+1}}{(\alpha-\beta)^{2m+2}}\right), \\ Q_2 = \frac{m(m+1)\alpha^{2m}}{(\alpha-\beta)^{2m+2}} + \frac{(m^2-1)\alpha^{2m-2}}{(\alpha-\beta)^{2m}} - \frac{2m\left(m+\frac{1}{2}\right)\alpha^{2m-1}}{(\alpha-\beta)^{2m+1}} \end{aligned}\right\} \tag{2.5}$$

Now the fundamental metric tensor $g_{ij} = \frac{1}{2}\dot{\partial}_i\,\dot{\partial}_j L^2$ is defined as,

$$g_{ij} = Pa_{ij} + P_0 b_i b_j + P_1\left(b_i Y_j + b_j Y_i\right) + Q_2 Y_i Y_j, \tag{2.6}$$

where,
$$P_0 = Q_0 + L_\beta^2, \qquad P_1 = Q_1 + L^{-1}PL_\beta, \qquad P_2 = Q_2 + P^2L^{-2}. \tag{2.7}$$

In addition, the reciprocal of tensor g_{ij} is g^{ij} given by

$$g^{ij} = P^{-1}a^{ij} - S_0 b^i b^j - S_1(b^i y^j + b^j y^i) - S_2 y^i y^j \tag{2.8}$$

where,

$$\left.\begin{aligned} b^i = a^{ij}b_j,\ S_0 = \frac{PP_0+(P_0P_2-P_1^2)\alpha^2}{\mu P},\\ S_1 = \frac{PP_1+(P_0P_2-P_1^2)\beta}{\mu P},\ S_2 = \frac{PP_2+(P_0P_2-P_1^2)b^2}{\mu P}\\ \mu = P(P+P_0b^2+P_1\beta)+(P_0P_2-P_1^2)(\alpha^2b^2-\beta^2). \end{aligned}\right\} \quad (2.9)$$

The Cartan tensor is given by,

$$2PC_{ijk} = P_1(h_{ij}m_k + h_{jk}m_i + h_{ki}m_j) + \gamma_1 m_i m_j m_k. \quad (2.10)$$

where,

$$C_{ijk} = \frac{1}{2}\dot{\partial}_k g_{ij}, \qquad \gamma_1 = P\left(\frac{\partial P_0}{\partial \beta}\right) - 3P_1Q_0, \qquad m_i = b_i - \alpha^{-2}\beta Y_i. \quad (2.11)$$

The associated Riemannian space containing the components of the Christoffel's symbol $\left\{ {i \atop jk} \right\}$ and the covariant derivative ∇_k with respect to x^k corresponding to this Christoffel's symbols.

We have the following components of the symmetric and skew symmetric tensors respectively,

$$E_{ij} = \frac{(b_{ij}+b_{ji})}{2}, \qquad F_{ij} = \frac{(b_{ij}-b_{ji})}{2}, \quad (2.12)$$

where $b_{ij} = \nabla_j b_i$.

The difference tensor $D^i_{jk} = \Gamma^{*i}_{jk} - \Gamma^i_{jk}$ of the special Finsler space F^n is given by

$$\begin{aligned} D^i_{jk} = B^i E_{jk} + F^i_k B_j + F^i_j B_k + B^i_j b_{0k} + B^i_k b_{0j} - b_{0m}g^{im}B_{jk} - C^i_{jm}A^m_k - C^i_{km}A^m_j + C_{jkm}A^m_s g^{is}\\ +\lambda^s(C^i_{jm}C^m_{sk} + C^i_{km}C^m_{sj} - C^m_{jk}C^i_{ms}) \end{aligned} \quad (2.13)$$

where,

$$\left.\begin{aligned} B_k = P_0 b_k + P_1 Y_k,\ B^i = g^{ij}B_j,\ F^k_i = g^{kj}F_{ji},\\ B_0 = b_i y^i,\ B_{ij} = \frac{\{P_1(a_{ij}-\alpha^{-2}Y_iY_j)+\frac{\partial P_0}{\partial \beta}m_i m_j\}}{2},\\ A^m_k = B^m_k E_{00} + B^m E_{k0} + B_k F^m_0 + B_0 F^m_k,\ B^k_i = g^{kj}B_{ji},\\ \lambda^m = B^m E_{00} + 2B_0 F^m_0. \end{aligned}\right\} \quad (2.14)$$

Here and also for the following we denote 0 as contraction with y^i except for the quantities P_0, Q_0 and S_0.

3. Induced Cartan Connection

Let F^{n-1} be a Hypersurface which contains an equation $x^i = x^i(u^\alpha), \alpha = 1,2,\dots(n-1)$, where u_α be a Gaussian coordinate on the hypersurface F^{n-1}. Let us consider that the matrix of the projection factor $B^i_\alpha = \frac{\partial x^i}{\partial u^\alpha}$ is of rank $(n-1)$, then

$$y^i = B^i_\alpha(u)v^\alpha \quad (3.1)$$

Here y^i is the supporting element of F^n is tangential to F^{n-1} and thus the $v = v^\alpha$ is the element of support of F^{n-1} at the point u^α. The metric tensor $g_{\alpha\beta}$ and Cartan tensor $C_{\alpha\beta\gamma}$ of F^{n-1} are given by

$$g_{\alpha\beta} = g_{ij}B^i_\alpha B^j_\beta, \qquad C_{\alpha\beta\gamma} = C_{ijk}B^i_\alpha B^j_\beta B^k_\gamma. \quad (3.2)$$

A unit normal vector $N^i(u,v)$ is at each point u_α of F^{n-1} is expressed by,

$$g_{ij}\big(x(u,v),y(u,v)\big)B^i_\alpha N^i = 0, \qquad g_{ij}\big(x(u,v),y(u,v)\big)N^iN^j = 1. \tag{3.3}$$

In terms of angular metric tensor h_{ij}, thus

$$h_{\alpha\beta} = h_{ij}B^i_\alpha B^j_\beta, \qquad h_{ij}B^i_\alpha N^j = 0, \qquad h_{ij}N^iN^j = 1. \tag{3.4}$$

If B^α_i is the inverse of B^i_α , then

$$\left.\begin{aligned} B^\alpha_i &= g^{\alpha\beta}g_{ij}B^j_\beta,\ B^i_\alpha B^\beta_i = \delta^\beta_\alpha,\ B^i_\alpha N^i = 0,\\ N_i &= g_{ij}N^j,\ B^i_\alpha B^\alpha_j + N^i N_j = \delta^i_j, \end{aligned}\right\} \tag{3.5}$$

where $g^{\alpha\beta}$ is the inverse of $g_{\alpha\beta}$ of F^{n-1}.

From Eq. (3.3) and Eq. (3.5), we have

$$B^i_\alpha B^\beta_i = \delta^\beta_\alpha, \qquad B^i_\alpha N_i = 0, \qquad N^iB^\alpha_i = 0, \qquad N^iN_i = 1. \tag{3.6}$$

And also, we have,

$$B^i_\alpha B^\alpha_j + N^iN_j = \delta^i_j. \tag{3.7}$$

The induced Cartan's connection $ICΓ = \left(\Gamma^{*\alpha}_{\ \beta\ \gamma}, G^\alpha_\beta, C^{\ \alpha}_{\beta\ \gamma}\right)$ of F^{n-1} generated from the Cartan's connection $CΓ = \left(\Gamma^{*i}_{\ j\ k}, \Gamma^{*i}_{0\ k}, C^{\ i}_{j\ k}\right)$ is given by,

$$\begin{aligned} \Gamma^{*\alpha}_{\ \beta\ \gamma} &= B^\alpha_i\left(B^i_{\beta\ \gamma} + \Gamma^{*i}_{\ j\ k}B^j_\beta B^k_\gamma\right) + M^\alpha_\beta H_\gamma\\ G^\alpha_\beta &= B^\alpha_i\left(B^i_{0\beta} + \Gamma^{*i}_{0\ j}B^j_\beta\right),\\ C^{\ \alpha}_{\beta\ \gamma} &= B^\alpha_i C^{\ i}_{j\ k}B^j_\beta B^k_\gamma. \end{aligned}$$

where,

$$M_{\beta\gamma} = N_iC^i_{jk}B^j_\beta B^k_\gamma, \quad M^\alpha_\beta = g^{\alpha\gamma}M_{\beta\gamma}, \quad H_\beta = \dot{N}_\iota\left(B^i_{0\beta} + \Gamma^{*i}_{0j}B^j_\beta\right)$$

and

$$B^i_{\beta\gamma} = \frac{\partial B^i_\beta}{\partial u^\gamma}, \quad B^i_{0\beta} = B^i_{\alpha\beta}v^\alpha$$

Here the second fundamental v-tensor and normal curvature vector are $M_{\beta\gamma}$ and H_β respectively, the second fundamental h-tensor $H_{\beta\gamma}$ is stated as [6]

$$H_{\beta\gamma} = N_i\left(B^i_{\beta\gamma} + \Gamma^{*i}_{jk}B^j_\beta B^k_\gamma\right) + M_\beta H_\gamma \tag{3.8}$$

where,

$$M_\beta = N_iC^i_{jk}B^j_\beta N^k n\,. \tag{3.9}$$

The second fundamental h-tensor and the normal curvature vector are $H_{\alpha\beta}$ and H_α of F^{n-1} respectively, are given by

$$H_{\alpha\beta} = N_i\left(B^i_{\alpha\beta} + F^i_{jk}B^j_\alpha B^k_\beta\right) + M_\alpha H_\beta \tag{3.10}$$

and

$$H_\alpha = N_i\left(B^i_{0\alpha} + G^i_j B^j_\alpha\right) \tag{3.11}$$

where $M_\alpha = C_{ijk}B^i_\alpha N^j N^k, B^i_{\alpha\beta} = \frac{\partial x^i}{\partial u^\alpha \partial u^\beta}$ and $B^i_{0\alpha} = B^i_{\beta\alpha}v^\beta$. It is clearly expresses that $H_{\alpha\beta}$ is not symmetric and

$$H_{\alpha\beta} - H_{\beta\alpha} = M_\alpha H_\beta - M_\beta H_\alpha. \tag{3.12}$$

The equations Eq. (3.10) and Eq. (3.11) gives

$$H_{0\alpha} = H_{\beta\alpha}v^\beta = H_\alpha, \quad H_{\alpha 0} = H_{\alpha\beta}v^\beta = H_\alpha + M_\alpha H_0. \tag{3.13}$$

Here,

$$M_{\alpha\beta} = C_{ijk}B^i_\alpha B^j_\beta N^k. \tag{3.14}$$

where $M_{\alpha\beta}$ is the second fundamental v-tensor. The projection factors B^α_i and N^i contains the h- and v-covariant derivatives evolve with respect to induced Cartan connection $IC\Gamma$ respectively are given by

$$B^i_{\alpha|\beta} = H_{\alpha\beta}N^i = M_{\alpha\beta}N^i, \quad N^i_{|\beta} = -H_{\alpha\beta}B^\alpha_j g^{ij} = -M_{\alpha\beta}B^\alpha_j g^{ij} \tag{3.15}$$

and in terms of h- and v-covariant derivatives of vector field X^i are as follows:

$$X_{i|\beta} = X_{i|j}B^j_\beta + X_{i|j}N^j H_\beta, \quad X_{i|\beta} = X_{i|j}B^j_\beta. \tag{3.16}$$

The following lemmas states the different kinds of hypersurfaces and their characteristics conditions which are defined by M. Matsumoto.

Lemma 3.1. *A hypersurface of a Finsler space F^{n-1} is a hyperplane of the 1^{st} kind if and only if $H_\alpha = 0$.*

Lemma 3.2. *A hypersurface of a Finsler space F^{n-1} is a hyperplane of the 2^{nd} kind if and only if $H_\alpha = 0$ and $H_{\alpha\beta} = 0$.*

Lemma 3.3. *A hypersurface of a Finsler space F^{n-1} is a hyperplane of the 3^{rd} kind if and only if $H_\alpha = 0$ and $M_{\alpha\beta} = H_{\alpha\beta} = 0$.*

4. Hypersurface F^{n-1} of Finsler space

Let us consider a Finsler space F^n with the (α, β)- metric $L = \frac{\alpha^{m+1}}{(\alpha-\beta)^m}$, where, $\alpha = \sqrt{a_{ij}(x)y^i y^j}$ is a Riemannian metric, $\beta = b_i(x)y^i$ is 1-form metric and $b_i(x) = \frac{\partial b}{\partial x^i}$ is a gradient of scalar function $b(x)$. In this part we prove the necessary and sufficient condition of hypersurface of a Finsler space to be hyperplane which satisfies the 1^{st} kind, 2^{nd} kind and but not 3^{rd} kind. We have,

$$b_i B^i_\alpha = 0, \quad b_i y^i = \beta = 0. \tag{4.1}$$

Accordingly, the induced metric $L(u, v)$ of hypersurface of Eq. (4.1) which is Riemannian metric given by

$$L(u, v) = \sqrt{a_{\alpha\beta}(u)v^\alpha v^\beta}, \quad a_{\alpha\beta} = a_{ij}B^i_\alpha B^j_\beta. \tag{4.2}$$

Henceforth $F^{n-1}(c)$, now from the equations Eq. (2.5), Eq. (2.7) and Eq. (2.9), we have

$$\left.\begin{aligned} P = 1,\ Q_0 = m(m+1),\ Q_1 = 0,\ Q_2 = -\tfrac{1}{\alpha^2},\ P_0 = m(2m+1), \\ P_1 = \tfrac{m}{\alpha},\ P_2 = 0,\ \mu = 1 + (m^2+m)b^2,\ S_0 = \tfrac{(m^2+m)}{1+(m^2+m)b^2}, \\ S_1 = \tfrac{m}{\alpha(1+(m^2+m)b^2)},\ S_2 = -\tfrac{m^2b^2}{\alpha^2(1+(m^2+m)b^2)}. \end{aligned}\right\} \tag{4.3}$$

wherefore Eq. (2.8) yields,

$$g^{ij} = a^{ij} - \frac{(m^2+m)}{1+(m^2+m)b^2} b^i b^j - \frac{m}{\alpha(1+(m^2+m)b^2)}(b^i y^j + b^j y^i) + \frac{m^2 b^2}{\alpha^2(1+(m^2+m)b^2)} y^i y^j. \quad (4.4)$$

Hence with F^{n-1}, operating the $b_i b_j$ using Eq. (4.1) and Eq. (4.4) we get

$$g^{ij} b_i b_j = \frac{b^2}{1+(m^2+m)b^2} \quad (4.5)$$

Therefore, we obtain

$$b_i\big(x(u)\big) = \sqrt{\frac{b^2}{1+(m^2+m)b^2}} N_i, \qquad b^2 = a^{ij} b_i b_j. \quad (4.6)$$

where b is the length of vector b^i. From Eq. (4.4) and Eq. (4.6), we get

$$b^i = a^{ij} b_j = \sqrt{b^2\big((m^2+m)b^2+1\big)} N^i + \frac{mb^2}{\alpha} y^i. \quad (4.7)$$

Theorem 4.1. *Let F^{n-1} be a hypersurface of Finsler space F^n with metric $L = \frac{\alpha^{m+1}}{(\alpha-\beta)^m}$ with scalar function $b_i(x) = \partial_i b(x)$ given by Eq. (4.5) and Eq. (4.6) and a hypersurface F^{n-1} of F^n. Then the induced Riemannian metric is led by (4.1).*

Theorem 4.2. *The second fundamental v-tensor of hypersurface of a Finsler space with the metric $L = \frac{\alpha^{m+1}}{(\alpha-\beta)^m}$ vanishes and the second fundamental h-tensor is symmetric.*

Proof: Further, h_{ij} and g_{ij} are identified by Eq. (2.3) and Eq. (2.6) into Eq. (4.3),

$$h_{ij} = a_{ij} + m(m+1) b_i b_j - \frac{1}{\alpha^2} Y_i Y_j \quad (4.8)$$

$$g_{ij} = a_{ij} + m(2m+1) b_i b_j + \frac{m}{\alpha}\big(b_i y_j + b_j y_i\big) \quad (4.9)$$

where, h_{ij} is an angular metric tensor and g_{ij} is a metric tensor.

Along hypersurface $F^{n-1}(c)$ with $h^{(a)}_{\alpha\beta}$ which represents the angular metric tensor of Riemannian $a_{ij}(x)$, then $h_{\alpha\beta} = h^{(a)}_{\alpha\beta}$. Accordingly, from Eq. (2.7), as well as $F^{n-1}(c)$, $\frac{\partial P_0}{\partial \beta} = 2m(m+1)(2m+1)\frac{\alpha^{2m+2}}{(\alpha-\beta)^{2m+3}}$.

Therefore, Eq. (2.11) yields $\gamma_1 = \frac{m(m+1)(m+2)}{\alpha}, m_i = b_i$.

Now from Eq. (2.10), we get

$$C_{ijk} = \frac{m}{2\alpha}\big(h_{ij} b_k + h_{jk} b_i + h_{ki} b_j\big) + \Big(\frac{m(m+1)(m+2)}{2\alpha}\Big) b_i b_j b_k. \quad (4.10)$$

Using Eq. (3.14), Eq. (3.4) and Eq. (4.10), we obtain

$$M_{\alpha\beta} = \frac{m}{2\alpha}\sqrt{\frac{b^2}{1+(m^2+m)b^2}} h_{\alpha\beta}, \quad M_\alpha = 0 \quad (4.11)$$

Thus from Eq. (4.11) and Eq. (3.12) thereby $H_{\alpha\beta} = H_{\beta\alpha}$, we have the following result:

Theorem 4.3. *The necessary and sufficient condition for a hypersurface $F^{n-1}(c)$ of a Finsler space with the generalized Matsumoto metric $L = \frac{\alpha^{m+1}}{(\alpha-\beta)^m}$ to be a hyperplane of first kind $b_{ij} = \frac{b_i c_j + b_j c_i}{2}$ holds.*

Proof: Subsequently, from Eq. (4.1) we get

$$b_{i|\beta} B^i_\alpha + b_i B^i_{\alpha|\beta} = 0. \quad (4.12)$$

Therefore from Eq. (3.15) and we are using Eq. (3.16), we get

$$b_{i|j}B^i_\alpha B^j_\beta + b_{i|j}B^i_\alpha N^j H_\beta + b_i H_{\alpha\beta}N^i = 0. \tag{4.13}$$

Along $b_{i|j} = -b_s C^s_{ij}$, becomes $b_{i|j}B^i_\alpha N^j = -\sqrt{\frac{b^2}{1+(m^2+m)b^2}}M_\alpha = 0$. Including $b_{i|j}$ is symmetric, from Eq. (4.13) we obtain

$$\sqrt{\frac{b^2}{1+(m^2+m)b^2}}H_{\alpha\beta} + b_{i|j}B^i_\alpha B^j_\beta = 0. \tag{4.14}$$

Now, contracting the Eq. (4.14) with v^β and again that with v^α, yields

$$\sqrt{\frac{b^2}{1+(m^2+m)b^2}}H_\alpha + b_{i|j}B^i_\alpha y^j = 0, \tag{4.15}$$

$$\sqrt{\frac{b^2}{1+(m^2+m)b^2}}H_0 + b_{i|j}y^i y^j = 0. \tag{4.16}$$

In the context of lemma 3.1, the hypersurface of Finsler space $F^{n-1}(c)$ is a hyperplane of the first kind if and only if $H_0 = 0$ and also on the other hand $b_{i|j}y^i y^j = 0$. The covariant derivative $b_{i|j}$ depends on y^i.

Since, from Eq. (2.12) which become $E_{ij} = b_{ij},\ F_{ij} = 0$ and $F^i_j = 0$.

Hence Eq. (2.13) reduces to

$$\begin{aligned} D^i_{jk} = {} & B^i\, b_{jk} + B^i_j\, b_{0k} + B^i_k b_{0j} - b_{0m}\, g^{im} B_{jk} C^i_{jm}\, A^m_k - C^i_{km}\, A^m_j \\ & + C_{jkm}\, A^m_s\, g^{is} + \lambda^s\left(C^i_{jm}\, C^m_{sk} + C^i_{km}\, C^m_{sj} - C^m_{jk} C^i_{ms}\right) \end{aligned} \tag{4.17}$$

In the context of Eq. (4.3) and Eq. (4.4), the relations in Eq. (2.14) consists of

$$\left.\begin{aligned} & B_i = m(2m+1)b_i + \tfrac{m}{\alpha}Y_i,\ \ B^i = \tfrac{m^2+m}{1+(m^2+m)b^2}b^i + \tfrac{m}{\alpha(1+(m^2+m)b^2)}y^i, \\ & \lambda^m = B^m\, b_{00},\ \ A^m_k = B^m_k\, b_{00} + B^m\, b_{k0}, B_{ij} = \tfrac{m}{2\alpha}a_{ij} - \tfrac{m}{2\alpha^3}Y_i Y_j + \tfrac{m(m+1)(2m+1)}{\alpha}b_i b_j, \\ & B^i_j = g^{ik}\, B_{kj} = \tfrac{m}{2\alpha}(\delta^i_j - \alpha^{-2}y^i Y_j) + \tfrac{m(m+1)(3m+2)}{2\alpha(1+(m^2+m)b^2)}b^i b_j - \tfrac{m^2+2m^2(m+1)(2m+1)b^2}{2\alpha^2(1+(m^2+m)b^2)}y^i b_j \end{aligned}\right\} \tag{4.18}$$

In the context of Eq. (4.3) and Eq. (4.4), the relations in Eq. (2.14) consists of

$$\left.\begin{aligned} D^i_{j0} &= B^i\, b_{j0} + B^i_j\, b_{00} - B^m\, C^i_{jm}\, b_{00}, \\ D^i_{00} &= B^i\, b_{00} = \left(\tfrac{m^2+m}{1+(m^2+m)b^2}b^i + \tfrac{m}{\alpha(1+(m^2+m)b^2)}y^i\right)b_{00}. \end{aligned}\right\} \tag{4.19}$$

By the relation Eq. (4.1), we obtain

$$b_i D^i_{j0} = \tfrac{(m^2+m)b^2}{1+(m^2+m)b^2}b_{j0} + \tfrac{m+3m(m+1)^2b^2}{2\alpha(1+(m^2+m)b^2)}b_j b_{00} - \tfrac{(m^2+m)b^m}{1+(m^2+m)b^2}b_i C^i_{jm} b_{00}. \tag{4.20}$$

$$b_i D^i_{00} = \tfrac{(m^2+m)b^2}{1+(m^2+m)b^2}b_{00}. \tag{4.21}$$

Hence $b_{i|j} = b_{ij} - b_s D^s_{ij}$ with Eq. (4.20) and Eq. (4.21) yields

$$b_{i|j}y^i y^j = \tfrac{1}{1+(m^2+m)b^2}b_{00}. \tag{4.22}$$

Subsequently Eq. (4.15) and Eq. (4.16) are expressed as

$$\tfrac{b}{\sqrt{1+(m^2+m)b^2}}H_\alpha + b_{i0}B^i_\alpha = 0. \tag{4.23}$$

$$\tfrac{b}{\sqrt{1+(m^2+m)b^2}}H_0 + \tfrac{1}{1+(m^2+m)b^2}b_{00} = 0. \tag{4.24}$$

As consequence of that the condition $H_0 = 0$ for an induced metric which is equivalent to $b_{00} = 0$, using Eq. (4.1) which can be written as $b_{ij}y^i y^j = (b_i y^i)(c_j y^j)$ for some $c_j(x)$, therefore

$$b_{ij} = \frac{b_i c_j + b_j c_i}{2}. \tag{4.25}$$

And also satisfies the hypersurface of the second kind, $b_{00} = 0,\ b_{ij}B^i_\alpha B^j_\beta = 0,\ b_{ij}B^i_\alpha y^j = 0.$

Here Eq. (4.24) gives $H_\alpha = 0$. And from Eq. (4.18) and Eq. (4.25) we obtain

$b_{i0}b^i = \frac{c_0 b^2}{2},\ \lambda^m = 0,\ A^i_j B^j_\beta = 0$ and $B_{ij}B^i_\alpha B^j_\beta = 0.$

With the help of Eq. (3.14), Eq. (4.4), Eq. (4.7), Eq. (4.11) and Eq. (4.17), we get

$$b_s D^s_{ij} B^i_\alpha B^j_\beta = -\frac{c_0 b^2 m}{4\alpha(1+(m^2+m)b^2)} h_{\alpha\beta},$$

henceforth, Eq. (4.14) reduces to

$$\frac{b}{\sqrt{1+(m^2+m)b^2}} H_{\alpha\beta} + \frac{c_0 b^2 m}{4\alpha(1+(m^2+m)b^2)} h_{\alpha\beta} = 0. \tag{4.26}$$

hence the result. From Eq. (3.13), Eq. (4.10), Eq. (4.18) and Eq. (4.16) and the Theorem 4.2.

Theorem 4.4. *The necessary and sufficient condition for a hypersurface $F^{n-1}(c)$ of a Finsler space with (α, β)-metric $L = \frac{\alpha^{m+1}}{(\alpha-\beta)^m}$ to be a hyperplane of second kind Eq. (4.26) holds. i.e., $H_{\alpha\beta} = 0$.*

Theorem 4.5. *The necessary and sufficient condition for a hypersurface $F^{n-1}(c)$ of a Finsler space with (α, β)-metric $L = \frac{\alpha^{m+1}}{(\alpha-\beta)^m}$ to be a hyperplane of third kind does not holds. we distinguished that the hypersurface $F^{n-1}(c)$ is not a hyperplane of the third kind.*

Here we concluded that the necessary and sufficient conditions for the hypersurface $F^{n-1}(c)$ of a Finsler space with the $L = \frac{\alpha^{m+1}}{(\alpha-\beta)^m}$, in this paper we determined the conditions for hypersrface to be a hyperplane of first, second kind and but not a third kind.

References

[1] M. Matsumoto, *Theory of Finsler spaces with (α, β)-metric*, Reports on mathematical physics, Vol. 31, 1992.

[2] K. Vineet and R. K. Gupta, *Hypersurface of a Special Finsler Space with Metric $L = \beta + \frac{\alpha^3+\beta^3}{\alpha(\alpha-\beta)}$*, Journal of Computer and Mathematical Sciences, Vol. 9, No. 6, pp. 579-587, June 2018.

[3] S. K. Narasimhamurthy, S. T. Aveesh, H. G. Nagaraja and Pradeep Kumar, *On special hypersurface of a Finsler space with the metric $\alpha + \frac{\beta^{n+1}}{\alpha^n}$*, Acta Universitatis Apulensis, Vol. 17, pp. 129-139, 2009.

[4] Pradeep Kumar, S. K. Narasimhamurthy, C. S. Bagewadi and S.T. Aveesh, *Special Finsler hypersurfaces admitting a parallel vector field*, SCIENTIA, Series A: Mathematical Science, Vol. 20, pp.123-130, 2010.

[5] U. P. Singh, and K. Bindu, *On a hypersurface of a Matsumoto space*, Indian J. pure appl. Math., Vol. 32, pp. 521-531, 2001.

[6] M. Matsumoto, *The induced and intrinsic Finsler connections of a hypersurface and Finslerien projective geometry*, Journal of Mathematics of Kyoto University, Vol. 25, No. 1, pp. 107-144, 1985.

[7] M. K. Gupta, Abhay Singh and P. N. Pandey, *On a Hypersurface of a Finsler space with Randers Change of Matsumoto metric*, Hindawi Publishing Corporation geometry, Article ID 842573, 6 pages, 2013.

[8] G. M. Brown, *The Study of Tensors which Characterize a Hypersurface of a Finsler space*, Canadian Journal of Mathematics, Vol. 20, pp. 1025 – 1036, 1968.

[9] H. Rund, *Hypersurface of a Finsler space*, Canadian Mathematical Congress, 1955.